

JOURNAL OF ALGEBRA 56, 401–408 (1979)

Endomorphism Rings and Lattice Isomorphisms

SOUMAYA MAKDISSI KHURI

*Yale University, New Haven, Connecticut 06520**Communicated by P. M. Cohn*

Received June 24, 1977

INTRODUCTION

A Baer ring is a ring in which every right (and left) annihilator ideal is generated by an idempotent. If B is a Baer ring, then the set of principal right ideals of B which are generated by idempotents forms a complete lattice which is anti-isomorphic to the lattice of principal left ideals generated by idempotents ([9], Proposition III 8.2). If ${}_R V$ is a vector space over a division ring, $\text{Hom}_R(V, V)$ is a Baer ring. In trying to determine whether $B = \text{Hom}_R(V, V)$ is a Baer ring for a general module ${}_R V$, a natural method to use is Baer's "Three-cornered Galois Theory," which establishes Galois connections between the lattice, $L(V)$, of submodules of V and the lattices $L(B_.)$ and $L(.B)$ of right and left ideals of B , and consequently lattice isomorphisms and anti-isomorphisms between a certain sublattice, \mathcal{C} , of $L(V)$, consisting of the "closed" submodules of V and the lattices \mathcal{A}_R and \mathcal{A}_L of right and left annihilators in B . This method has been used in [11] and [10] for the case of a free module and in [6] for the case of a projective module which contains a unimodular element.

In both the free and the projective cases, two key steps are needed in order to establish the lattice isomorphisms: the first step consists in proving that the module V is a self-generator (i.e. $V \text{Hom}_R(V, U) = U$ for every submodule U of V), while the second step consists in proving that the class of "closed" submodules under consideration (which, in [6], [10] and [11] happen to be the "dual-closed" submodules, i.e. the submodules which equal their own second annihilators with respect to the dual space $V^* = \text{Hom}_R(V, R)$ of V) coincides with the class, \mathcal{C}_a , of "annihilator-closed" submodules of V , where the annihilator-closure operator is the one obtained from the Galois connection between $L(V)$ and $L(B_.)$ which is given by Baer's three-cornered Galois Theory.

In the absence of "free-ness" or of a unimodular element, the module V need not be a self-generator; however, we show, in Section 2, that an "approximation" of self-generation is sufficient for the lattice (anti-) isomorphism between $(\mathcal{A}_R) \mathcal{A}_L$ and \mathcal{C}_a to hold; and this allows us to escape from the "free" and the

“unimodular” situations. This approximation, which we call c -self-generation (where $\varphi(U) = U^c$ is the closure operator under consideration), consists in the module V satisfying $[V \operatorname{Hom}_R(V, U)]^c = U$, for each c -closed submodule U . Two important and frequently considered closure operators are: the essential-closure operator, which we shall denote by $\varphi(U) = U^e$, and the dual-closure operator, $\varphi(U) = U^d$. We show, in Section 3, that any nonsingular module ${}_R V$, with the essential-closure operator, is an e -self-generating module, provided it satisfies the rather mild condition that $\operatorname{Hom}_R(V, U) \neq 0$ for every nonzero essentially-closed submodule U ; and that any module ${}_R V$ with the dual-closure operator is d -self-generating, provided that V^* is T -faithful, where T is the trace (V, V^*) of V and we say, following Sandomierski ([8]), that a module X is T -faithful if $Xt \neq 0$ for each nonzero $t \in T$.

Whereas the first step of the method mentioned above can be generalized to modules which are not necessarily free or projective by requiring the modules to be c -self-generating, the second step must be resolved individually for each particular type of submodule and its closure operator. For a nonsingular module with the essential-closure operator, Theorem 3.5 gives necessary and sufficient conditions for the essentially closed submodules to coincide with the annihilator-closed ones, while for a general module with dual-closure, a sufficient condition is given in Proposition 3.6 for the dual-closed submodules to coincide with the annihilator-closed ones. Finally, applications of the three-cornered Galois Theory to Baer endomorphism rings are given in Theorems 3.7 and 3.8.

2. THE LATTICE ISOMORPHISMS

Throughout this paper, R denotes an associative ring with 1, ${}_R V$ a unitary left R -module and $B = \operatorname{Hom}_R(V, V)$. The action of elements of B on V will be written on the right. The right (left) annihilator in B of a subset H of B will be denoted by $\mathcal{R}(H)$ ($\mathcal{L}(H)$), while r and l will be used for annihilators in V of subsets of B , or in B of subsets of V , e.g. $l_V(H) = \{v \in V: vh = 0, \forall h \in H\}$ and $r_B(U) = \{b \in B: ub = 0 \forall u \in U\}$, $U \subseteq V$. Also, let $I(U) = \{b \in B: Vb \subseteq U\}$ and $UH = \{uh: u \in U \text{ and } h \in H\}$.

The following lemma is straightforward ([11], Lemma 1.1):

LEMMA 2.1. *If $U \subseteq V$ and $J \subseteq B$, then*

- (i) $VI(U) \subseteq U$.
- (ii) $U \subseteq l_V r_B(U)$.
- (iii) $I(U) r_B(U) = 0$.
- (iv) $Il_V(J) = \mathcal{L}(J)$.
- (v) $r_B(VJ) = \mathcal{R}(J)$. ■

Let L be a complete lattice. A *closure operator* on L is a mapping $\varphi: L \rightarrow L$, written $\varphi(a) = a^c$, such that:

- (c1) $a \leq b$ implies $a^c \leq b^c$;
- (c2) $a \leq a^c$;
- (c3) $(a^c)^c = a^c$.

An element a is *c-closed* if $a = a^c$. Given a closure operator, $\varphi(a) = a^c$, on a lattice L , we denote by \mathcal{C}_c the set of *c-closed* elements of L .

Let L' be another complete lattice. A *Galois connection* between L and L' is a pair of mappings $\sigma: L \rightarrow L'$ and $\tau: L' \rightarrow L$ satisfying:

- (1) $x_1 \leq x_2$ implies $\sigma(x_1) \geq \sigma(x_2)$ for $x_1, x_2 \in L$.
- (2) $y_1 \leq y_2$ implies $\tau(y_1) \geq \tau(y_2)$ for $y_1, y_2 \in L'$.
- (3) $x \leq \tau\sigma(x)$ and $y \leq \sigma\tau(y)$ for $x \in L, y \in L'$.

Given a Galois connection (σ, τ) , it can be shown that $\sigma\tau\sigma(x) = \sigma(x)$ and $\tau\sigma\tau(y) = \tau(y)$ for $x \in L, y \in L'$, so that the maps $\tau\sigma$ and $\sigma\tau$ are closure operators on L and L' , respectively. The closed elements in L are those which are of the form $\tau(y)$ for some $y \in L'$. σ and τ induce an anti-isomorphism between the corresponding lattices of closed elements ([9], pp. 76-78).

It is easily seen that \mathcal{L} and \mathcal{R} form a Galois connection between the lattice $L(B.)$ of right ideals of B and the lattice $L(.B)$ of left ideals of B , and that the mappings r_B and l_V form a Galois connection between the lattice $L(V)$ of submodules of V and the lattice $L(B.)$, giving the closure operators $r_B l_V$ and $l_V r_B$ on $L(V)$ and $L(B.)$ respectively. Let $\mathcal{C}_a = \{U \subseteq V: U = l_V r_B(U)\}$ be the collection of closed submodules of V with respect to the annihilator-closure operator $\varphi(U) = l_V r_B(U) = U^a$; the members of \mathcal{C}_a will be referred to as the "annihilator-closed" or "*a-closed*" submodules of V .

Sandomierski ([7]) has called a module ${}_R V$ a *self-generator* if $VI(U) = U$ for every submodule U of V . In general, one only has $VI(U) \subset U$ (cf. Lemma 2.1 (i)); thus, in the presence of a closure operator, $\varphi(U) = U^c$, on $L(V)$, it is natural to approximate self-generation by requiring that $VI(U)$ and U have the same closure, at least when U is *c-closed*.

DEFINITION 1. Let ${}_R V$ be a module with a closure operator, $\varphi(U) = U^c$, defined on its lattice of submodules, $L(V)$. V will be called a *c-self-generator* if $[VI(U)]^c = U$ for each *c-closed* $U \in L(V)$.

DEFINITION 2. Let ${}_R V$ be a module with a closure operator, $\varphi(U) = U^c$. A subring D of $B = \text{Hom}_R(V, V)$ will be called *c-continuous* if

$$Xf \subseteq (Xf)^c \quad \forall f \in D \quad \text{and} \quad X \in \mathcal{L}(V).$$

We shall show in the next section that any subring of B for a module (resp. a nonsingular module) with dual-closure (resp. essential-closure) is d -continuous (resp. e -continuous).

LEMMA 2.2. *Let ${}_R V$ be a module with a closure operator, $\varphi(U) = U^c$; if $B = \text{Hom}_R(V, V)$ is c -continuous then: (i) $r_B(X) = r_B(X^c)$ for $X \in \mathcal{L}(V)$, and (ii) $l_V(J) = [l_V(J)]^c$ for $J \subseteq B$.*

Proof. Clearly, $X \subseteq X^c$ implies $r_B(X^c) \subseteq r_B(X)$; on the other hand, if $b \in r_B(X)$, then $X^c b \subseteq (Xb)^c = 0$, hence $b \in r_B(X^c)$, proving equality. Now let $U = l_V(J)$; if $x \in U^c$, then $xJ \subseteq U^c J \subseteq (UJ)^c = 0$, therefore $x \in l_V(J) = U$, i.e. $U^c \subseteq U$, hence $U = U^c$. ■

Remark. Since each member of \mathcal{C}_a may be written in the form $l_V(J)$, for $J \subseteq B$, Lemma 2.2 (ii) implies that, for B c -continuous, $\mathcal{C}_a \subseteq \mathcal{C}_c$.

THEOREM 2.3. *Let ${}_R V$ be a c -self-generator and B be c -continuous.*

- (a) *If $U = U^c$, then $r_B(U) = \mathcal{R}[I(U)]$.*
- (b) *If $J = \mathcal{R}(H)$, for $H \subseteq B$, then $J = r_B l_V(J)$.*
- (c) *If $H = \mathcal{L}(J)$, for $J \subseteq B$, then $l_V(J) = (VH)^c$ and $H = I[(VH)^c]$.*

Proof. (a) Let $U = U^c$; then $r_B(U) = r_B[(VI(U))^c] = r_B[VI(U)]$ (by Lemma 2.2 (i)) $= \mathcal{R}[I(U)]$ (by Lemma 2.1 (v)).

(b) Let $J = \mathcal{R}(H)$, for $H \subseteq B$, and set $U = l_V(J)$. By Lemma 2.2 (ii), $U = U^c$, hence, by (a), $r_B(U) = \mathcal{R}[I(U)]$, or $r_B l_V(J) = \mathcal{R}[Il_V(J)] = \mathcal{R}[\mathcal{L}(J)]$ (by Lemma 2.1 (iv)) $= \mathcal{R}\mathcal{L}\mathcal{R}(H) = \mathcal{R}(H) = J$.

(c) Let $H = \mathcal{L}(J)$, where $J \subseteq B$, and set $U = l_V(J)$. Since $U = U^c$ and V is a c -self-generator, $U = [VI(U)]^c$, or $l_V(J) = [VI l_V(J)]^c = [V\mathcal{L}(J)]^c = (VH)^c$. Consequently, $I[(VH)^c] = Il_V(J) = \mathcal{L}(J) = H$. ■

Let

$$\begin{aligned} \mathcal{A}_{\mathcal{R}} &= \{J \subseteq B: J \text{ is a right annihilator}\} & \text{and} \\ \mathcal{A}_{\mathcal{L}} &= \{H \subseteq B: H \text{ is a left annihilator}\}. \end{aligned}$$

THEOREM 2.4. *If ${}_R V$ is a c -self-generator and B is c -continuous then: (a) The maps $\mathcal{A}_{\mathcal{R}} \rightarrow \mathcal{C}_a$ by $J \rightarrow l_V(J)$ and $\mathcal{C}_a \rightarrow \mathcal{A}_{\mathcal{R}}$ by $U \rightarrow r_B(U)$ determine a lattice anti-isomorphism between \mathcal{C}_a and $\mathcal{A}_{\mathcal{R}}$. (b) The maps $\mathcal{A}_{\mathcal{L}} \rightarrow \mathcal{C}_a$ by $H \rightarrow (VH)^c$ and $\mathcal{C}_a \rightarrow \mathcal{A}_{\mathcal{L}}$ by $U \rightarrow I(U)$ determine a lattice isomorphism between \mathcal{C}_a and $\mathcal{A}_{\mathcal{L}}$.*

Proof. (a) Let $U \in \mathcal{C}_a$, then, since $\mathcal{C}_a \subseteq \mathcal{C}_c$, Theorem 2.3 (a) applies and $r_B(U) = \mathcal{R}[I(U)]$, i.e. $r_B(U) \in \mathcal{A}_{\mathcal{R}}$. Let $J \in \mathcal{A}_{\mathcal{R}}$, then, clearly, $l_V r_B[l_V(J)] = l_V(J)$, so that $l_V(J) \in \mathcal{C}_a$. Since, by Theorem 2.3 (b), $J \in \mathcal{A}_{\mathcal{R}}$ implies $J = r_B l_V(J)$, and, by definition, $U \in \mathcal{C}_a$ implies $U = l_V r_B(U)$, it is now clear that r_B and l_V determine a lattice anti-isomorphism between \mathcal{C}_a and $\mathcal{A}_{\mathcal{R}}$.

(b) If $U = l_V r_B(U)$ is in \mathcal{C}_a , then, using Lemma 2.1 (iv), $l(U) = \Pi_V r_B(U) = \mathcal{L}[r_B(U)] \in \mathcal{A}_{\mathcal{L}}$. Let $H = \mathcal{L}(J) \in \mathcal{A}_{\mathcal{L}}$; then, by Theorem 2.3 (c), $(VH)^c = l_V(J)$ is in \mathcal{C}_a .

Again by Theorem 2.3 (c), $H \in \mathcal{A}_{\mathcal{L}}$ implies $\Pi[(VH)^c] = H$, and $U \in \mathcal{C}_a$ implies $[VI(U)]^c = [V\mathcal{L}(r_B(U))]^c = l_V r_B(U) = U$, showing that the given maps determine a lattice isomorphism. ■

3. EXAMPLES

A module ${}_R V$ is an *essential extension* of a submodule U —written $U \subset' V$ —if every nonzero submodule of V has nonzero intersection with U . One then says that U is *essential* in V . A submodule U of a module V is said to be *essentially closed* in V if U has no proper essential extensions in V . For $v \in V$, set $[U: v] = \{r \in R: rv \in U\}$; it is known that, if $U \subset' V$, then, for any nonzero $v \in V$, $[U: v] \subset' R$. The *singular submodule* $Z_R(V)$ of V is defined to be $\{v \in V: [0: v] \subset' R\}$. V is said to be *nonsingular* if $Z_R(V) = 0$. If $U \subset' V$, then V is nonsingular if and only if U is nonsingular. A ring will be called (left) nonsingular if its left regular representation is nonsingular. For details on essential extensions and nonsingular modules see [2] or [3]. If ${}_R V$ is a nonsingular module, then each submodule U of V has a unique maximal essential extension U^e in V , called the *essential closure* of U in V . This defines a closure operator, $\varphi(U) = U^e$, on $L(V)$.

PROPOSITION 3.1. *If ${}_R V$ is a nonsingular module and $\varphi(U) = U^e$ is the essential closure of a submodule U , then $B = \text{Hom}_R(V, V)$ is e -continuous.*

Proof. This is Lemma 3.2 of [5].

A module ${}_R V$ has been called *compressible* if, for each nonzero submodule X of V , $\text{Hom}_R(V, X)$ contains a nonzero monomorphism ([1]). Generalizing this concept, we will call a module ${}_R V$ *retractable* if $\text{Hom}_R(V, U) \neq 0$ for each nonzero submodule U of V . If ${}_R V$ has a closure operator, $\varphi(X) = X^c$, then ${}_R V$ will be called *c-retractable* if $\text{Hom}_R(V, U) \neq 0$ for each nonzero c -closed submodule U . For example, any torsionless module over a semiprime ring is retractable (cf. [12] Proposition 1.2 (ii)).

PROPOSITION 3.2. *If ${}_R V$ is a nonsingular e -retractable module, then V is e -self-generating.*

Proof. This proved in [5] (Theorem 3.4: (ii) \Leftrightarrow (iii)).

Consider now a module ${}_R V$ in its standard Morita context, (R, V, V^*, B) , that is, $V^* = \text{Hom}_R(V, R)$, $B = \text{Hom}_R(V, V)$, and ${}_R V_B$ and ${}_B V_R^*$ are, in a natural way, bimodules with an $R - R$ bimodule homomorphism $(\ , \) : V \otimes_B V^* \rightarrow R$ and a $B - B$ bimodule homomorphism $[\ , \] : V^* \otimes_R V \rightarrow B$ given

by $(v, g = vg)$ (the evaluation map) and $(\cdot)[g, v] = (\cdot, g)v$ for $v \in V$ and $g \in V^*$; so that (\cdot, \cdot) and $[\cdot, \cdot]$ satisfy:

$$v_1[g_1, v_2] = (v_1, g_1)v_2 \text{ and } g_1(v_1, g_2) = [g_1, v_1]g_2, \forall v_1, v_2 \in V, g_1, g_2 \in V^*.$$

Note also that $(vb, g) = (v, bg)$ and $[gr, v] = [g, rv] \forall v \in V, b \in B, g \in V^*$ and $r \in R$. For $X \subseteq V$, define the annihilator of X in V^* by $X^\perp = \{g \in V^* : (X, g) = 0\}$, and for $Z \subseteq V^*$, define $Z = \{v \in V : (v, Z) = 0\}$. A submodule U of V will be called *dual-closed* if $U = {}^\perp U^\perp$. Clearly, $U \subseteq {}^\perp U^\perp$, ${}^\perp U^{\perp\perp} = U^\perp$ and $\varphi(U) = U^d = {}^\perp U^\perp$ defines a closure operator on $L(V)$.

PROPOSITION 3.3. *For any module ${}_R V$ with dual-closure, $\varphi(U) = U^d$, $B = \text{Hom}_R(V, V)$ is d -continuous.*

Proof. To show continuity, we need to show $(X^d)b \subseteq (Xb)^d, \forall b \in B$. Let $y \in X^d = {}^\perp X^\perp$, so that $(y, X^\perp) = 0$; to show $yb \in (Xb)^d = {}^\perp (Xb)^\perp$, we must show $(yb, (Xb)^\perp) = 0$. Let $g \in (Xb)^\perp$, so that $(Xb, g) = 0$; then $(X, bg) = 0$, i.e. $bg \in X^\perp$. But this last implies $(y, bg) = 0$ and therefore $(yb, g) = 0$, as required. ■

For a module ${}_R V$ with trace $T = (V, V^*)$, T was called V -faithful in [8] if $Tv \neq 0$ for each $0 \neq v \in V$. Analogously, we shall say that V^* is T -faithful if $V^*t \neq 0$ for each $0 \neq t \in T$.

PROPOSITION 3.4. *If V^* is T -faithful, then V is d -self-generating.*

Proof. Since $VI(U) \subseteq U$ implies $U^\perp \subseteq [VI(U)]^\perp$, it suffices to prove $[VI(U)]^\perp \subseteq U^\perp$. Let $g \in [VI(U)]^\perp$, i.e. $(VI(U), g) = 0$. Then, for any $b \in I(U)$, $(V, bg) = (Vb, g) = 0$, and therefore $bg = 0$. This means $I(U)g = 0 \forall g \in [VI(U)]^\perp$. Now, for any $u \in U$, $v[V^*, u] = (v, V^*)u \subseteq Ru$, i.e. $[V^*, u] \subseteq I(U)$, hence, by the preceding, $[V^*, u]g = 0 \forall g \in [VI(U)]^\perp$, or $V^*(u, g) = 0$. But since V^* is T -faithful, this implies $(u, g) = 0$, i.e. $(U, g) = 0$ for $g \in [VI(U)]^\perp$, or $g \in U^\perp$. ■

We consider now the question of when the a -closure of a submodule coincides with its e -closure or with its d -closure. In the case of a nonsingular module with essential-closure, we have a complete answer in the following, where $E(V)$ denotes the injective hull of V .

THEOREM 3.5. *Let ${}_R V$ be a nonsingular module and let $\varphi(U) = U^e$ be the essential closure of a submodule U of V . Then the following are equivalent:*

- (i) $U^e = U^a$ for every submodule U of V .
- (ii) $r_B(U) \neq 0$ for every $V \neq U \in \mathcal{C}_e$.
- (iii) Every nonzero right ideal of $\text{Hom}_R(E(V), E(V))$ has nonzero intersection with $B = \text{Hom}_R(V, V)$.
- (iv) $U = U^e \Rightarrow U = l_V(J)$, for some subset J of B .

This is Theorem 3.5 of [5]. An example of a module satisfying the equivalent conditions of Theorem 3.5 is given by any finite-dimensional torsionless module over a ring which possesses a two-sided quotient ring ([4], Theorem 3.5).

For dual-closure, the following result holds:

PROPOSITION 3.6. *If ${}_R V$ is a T -faithful module, where $T = (V, V^*)$ is the trace of V , then ${}^\perp U^\perp = U^a$ for each submodule U of V .*

Proof. By d -continuity of $B = \text{Hom}_R(V, V)$ (Proposition 3.3), $U = l_V(J)$ is dual-closed, so that ${}^\perp U^\perp \subseteq U^a$. Conversely, let $x \in U^a = l_V r_B(U)$. If $x \notin {}^\perp U^\perp$, then there is $g \in U^\perp$ such that $(x, g) \neq 0$. Since V is T -faithful, $(x, g) V \neq 0$, i.e. there is $0 \neq v \in V$ such that $(x, g) v = 0$. Consider $[g, v] = b \in B$: we have $Ub = U[g, v] = (U, g) v = 0$, so that $b \in r_B(U)$, and yet $xb = x[g, v] = (x, g) v \neq 0$, contradicting $x \in l_V r_B(U)$. Hence $x \in {}^\perp U^\perp$ and $U^a = {}^\perp U^\perp$. ■

Note that any ${}_R V$, where R is semiprime, is T -faithful: given $0 \neq t = (y, f) \in T$, the semiprime-ness of R gives $0 \neq r \in R$ such that $(y, f) r(y, f) \neq 0$, and hence, in particular, $(y, f) r y \neq 0$. In fact, when R is semiprime, V^* is T -faithful also: if $0 \neq (y, f) \in T$, then $(y, f) r(y, f) \neq 0$, for $0 \neq r \in R$, also implies $f r(y, f) \neq 0$. Hence, for a module ${}_R V$ over a semiprime ring, there is a lattice (anti-)isomorphism between the (right) left annihilators in B and the dual-closed submodules of V .

Also, if V is T -faithful and T is V -faithful then V^* is T -faithful:

Proof.

$$\begin{aligned} V^*(v, w) = 0 &\Rightarrow [V^*(v, w), V] = 0 \Rightarrow [V^*, (v, w) V] = 0, \\ &\Rightarrow V[V^*, (v, w) V] = 0 \Rightarrow (V, V^*)(v, w) V = 0 \\ &\Rightarrow (v, w) V = 0, \quad \text{since } T \text{ is } V\text{-faithful} \\ &\Rightarrow (v, w) = 0, \quad \text{since } V \text{ is } T\text{-faithful.} \end{aligned}$$

When V is B -faithful, as is the case here, the condition “ T is V -faithful” is equivalent to the “non-degeneracy condition: $[V^*, v] \neq 0$ for $v \neq 0$ ”. Hence, in particular, if V is an R -faithful module satisfying the non-degeneracy condition, then there is a lattice (anti-)isomorphism between the (right) left annihilators in B and the dual-closed submodules of V .

A straightforward application of the methods and proofs of [11] yields the following:

THEOREM 3.7. *If R is semiprime or if ${}_R V$ is faithful and satisfies the non-degeneracy condition, then $B = \text{Hom}_R(V, V)$ is a Baer ring if and only if every dual-closed submodule of V is a direct summand in V .*

THEOREM 3.8. *If ${}_R V$ is a finite-dimensional torsionless module over a ring R possessing a two-sided quotient ring, B is a Baer ring if and only if every essentially closed submodule of V is a direct summand in V .*

ACKNOWLEDGMENT

The author is greatly indebted and deeply grateful to Professor Charles E. Rickart for many helpful discussions and suggestions which substantially improved the scope of this paper.

REFERENCES

1. A. K. BOYLE, Injectives containing no proper quasi-injective submodules, *Comm. Alg.* **4** (1976), 775–785.
2. C. FAITH, “Lectures on Injective Modules and Quotient Rings,” Lecture Notes in Mathematics No. 246, Springer-Verlag, New York, 1972.
3. K. R. GOODEARL, “Ring Theory,” Dekker, New York, 1976.
4. S. M. KHURI, “Baer Rings of Endomorphisms,” Ph. D. Dissertation, Yale University, 1974.
5. S. M. KHURI, Baer endomorphism rings and closure operators, *Can. J. Math.*, Vol. XXX, No. 5 (1978), 1070–1078.
6. B. R. McDONALD, Endomorphism rings of infinitely generated projective modules, *J. Algebra* **45** (1977), 69–82.
7. F. L. SANDOMIERSKI, “Relative Injectivity and Projectivity,” Ph. D. Dissertation, Pennsylvania State University, 1964.
8. F. L. SANDOMIERSKI, Modules over the endomorphism ring of a finitely generated projective module, *Proc. Amer. Math. Soc.* **31** (1972), 27–31.
9. B. STENSTRÖM, “Rings of Quotients,” Grundlehren der mathematischen Wissenschaften, Vol. 217, Springer-Verlag, Berlin/New York, 1975.
10. G. M. TSUKERMAN, Rings of endomorphisms of a free module, *Siberian Mat. Z.* **7** (1966), 1161–1167 (Russian); translated in *Siberian Math. J.* **7** (1966), 923–927.
11. K. G. WOLFSON, Baer rings of endomorphisms, *Math. Ann.* **143** (1961), 19–28.
12. J. M. ZELMANOWITZ, Endomorphism rings of torsionless modules, *J. Algebra* **5** (1967), 325–341.